

Broadleaf Weed Control with Status[®] Herbicide

Technical Information Bulletin



- ♥ Rapid Knockdown
- ♥ Residual Control
- ♥ Resistance Management

Rapid Knockdown

Effect of Status[®] herbicide (2.5 oz/A) on Lambsquarters and Morningglory within Hours After Treatment

3 Hours-After-Treatment

20 Hours-After-Treatment



Untreated

Roundup[®]
alone

Status +
Roundup

Untreated

Roundup[®]
alone

Status +
Roundup

Additional Features and Benefits:

- **Broad-spectrum Weed Control:** Excellent control of over 65 broadleaf weeds
- **Crop safe:** Contains a patented crop safener providing excellent crop safety
- **Simple to use:** Excellent stand-alone herbicide and tank-mix partner with Roundup in Roundup Ready[®] corn

Broad-spectrum Weed Control

Weeds	Status [®]	Callisto [®]	Laudis [™]
Amaranth spp.	C	C	C
Buckwheat, wild	C	PC	PC
Cocklebur, common	C	C	C
Dock, curly	C	PC	PC
Lambsquarters, common	C	C	C
Marestail / Horseweed	C	PC	PC
Morningglory spp.	C	PC	PC
Pigweed spp.	C	C	C
Pusley, Florida	C	C ¹	C ¹
Ragweed spp.	C	PC ²	C
Sicklepod	C	NC	PC
Sida, prickly (teaweed)	C	NC	NC
Smartweed spp.	C	C	C
Velvetleaf	C	C	C
Waterhemp	C	C	C
Atrazine recommended for improved weed control	No	Yes	Yes
Crop rotational restrictions	No³	Yes	Yes

C = Control; PC = Partial Control; NC = No Control

¹Control if applied to weeds <2 inches in height.

²Giant ragweed = C; Common ragweed = PC

³Following at least 1 inch of rainfall or overhead irrigation after the last application of Status